

MYCOTAXON

<http://dx.doi.org/10.5248/130.465>

Volume 130, pp. 465–469

April–June 2015

***Cryptocoryneum parvulum*, a new species
on *Araucaria angustifolia* (Brazilian pine)**

SILVANA SANTOS DA SILVA¹, LUÍS FERNANDO PASCHOLATI GUSMÃO^{1*},
& RAFAEL F. CASTAÑEDA-RUIZ²

¹*Departamento de Ciências Biológicas, Laboratório de Micologia, Universidade Estadual
de Feira de Santana, BR116 KM 03, 44031-460, Feira de Santana, Brazil*

²*Instituto de Investigaciones Fundamentales en Agricultura Tropical ‘Alejandro de Humboldt’ (INIFAT),
Académico Titular de la Academia de Ciencias de Cuba,
Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200*

*CORRESPONDENCE TO: lgusmao@uefs.br

ABSTRACT—*Cryptocoryneum parvulum* sp. nov., collected on decaying needle-like leaves of *Araucaria angustifolia*, is described and illustrated. It is distinguished by clavate inflated conidiophores and small cheiroid conidia with black cap cells and four pendulous arms of pale brown cells. Two other *Cryptocoryneum* species, *C. condensatum* and *C. rilstonei*, are newly recorded and illustrated. A key to *Cryptocoryneum* species found in Brazil is provided.

KEY WORDS— asexual fungi, systematics, subtropical fungi

Introduction

During a mycological survey of fungi associated with litter at a Brazilian araucaria forest, an interesting *Cryptocoryneum* specimen was collected from decaying needle-like leaves. It showed remarkable differences from all previously described *Cryptocoryneum* species (Ellis 1971, 1976) and is therefore described here as new.

Materials & methods

Samples of litter were placed in paper bags, taken to the laboratory, and prepared according to Castañeda-Ruiz (2005). Mounts were prepared in PVL (polyvinyl alcohol, lactic acid, and phenol) and measurements were made at a magnification of ×1000. Micrographs were obtained with an Olympus BX51 microscope equipped with bright field and Nomarski interference optics. The type specimen is deposited in the Herbarium of Universidade Estadual de Feira de Santana (HUEFS).

Taxonomy

Cryptocoryneum parvulum S.S. Silva, Gusmão & R.F. Castañeda, **sp. nov.** FIG. 1
IF 550828

Differs from other *Cryptocoryneum* spp. by its inflated conidiophores and conidiogenous cells and its smaller, cheiroid conidia with dark cap cells and four pendulous, 3–4-septate, pale brown to subhyaline arms.

TYPE: Brazil, Rio Grande do Sul State: São Francisco de Paula, Floresta Nacional de São Francisco de Paula, 29°25 S 50°23 W, 838 m alt., on decaying needle-like leaves of *Araucaria angustifolia* (Bertol.) Kuntze (*Araucariaceae*), 27.VIII.2014, coll. S.S. Silva. (Holotype, HUEFS 210439).

ETYMOLOGY: Latin, *parvulum*, referring to the small conidia.

CONIDIOMATA sporodochial, pulvinate, 75–140 µm diam., scattered, black. Mycelium superficial and immersed. Hyphae septate, branched, 2.5–3 µm diam., smooth, brown. CONIDIOPHORES distinct, clavate, inflated, single, erect, straight, close packet, pale brown, 0–2-septate, smooth, 15–25 × 9–16.5 µm. CONIDIOGENOUS CELLS monoblastic, clavate to cylindrical or doliiform, integrated, terminal, determinate. CONIDIA solitary, acrogenous, cheiroid, branched, 12–23 × 10.5–15 µm, with black cap cells and four pendulous pale brown arms, smooth, 3–4-septate, 2.5–5 µm wide.

NOTE: *Cryptocoryneum* Fuckel (Ellis 1971, 1976, Schoknecht & Crane 1977) is characterized by monoblastic conidiogenous cells, and cheiroid conidia composed of a series of dark brown cap cells from which the pendulous arms develop.

Additional *Cryptocoryneum* records from Brazil

Cryptocoryneum condensatum (Wallr.) E.W. Mason & S. Hughes ex S. Hughes,
Can. J. Bot. 36: 758 (1958). FIG. 2 A–D

SPECIMEN EXAMINED: BRAZIL, RIO GRANDE DO SUL STATE, São Francisco de Paula, Floresta Nacional de São Francisco de Paula, 29°25 S 50°23 W, 838 m alt., on decaying needle-like leaves of *Araucaria angustifolia*, 29.IX.2014, coll. S.S. Silva (HUEFS 210440).

NOTE: A new record for Brazil.

Cryptocoryneum rilstonei M.B. Ellis, Mycol. Pap. 131: 2 (1972). FIG. 2 E–H

SPECIMEN EXAMINED: BRAZIL, RIO GRANDE DO SUL STATE, São Francisco de Paula, Floresta Nacional de São Francisco de Paula 29°25 S 50°23 W, 838 m alt., on decaying needle-like leaves of *Araucaria angustifolia*, 21.VIII.2014, coll. S.S. Silva (HUEFS 210441).

NOTE: A new record for Brazil.




FIG. 1. *Cryptocoryneum parvulum* (holotype, HUEFS 210439): A–B. Sporodochia on natural substrate. C–E. Sporodochia. F. Conidiophores, conidiogenous cells and conidia. G–H. Conidiogenous cell and conidia. I. Conidia.



FIG 2. *Cryptocoryneum condensatum* (HUEFS 210440): A. Sporodochium on the natural substratum. B–D. Conidia. *Cryptocoryneum rilstonei* (HUEFS 210441): E. Sporodochium on the natural substratum. F. Sporodochium. G–H. Conidia.

Key to *Cryptocoryneum* species found in Brazil

1. Conidiophores cylindrical 2
1. Conidiophores clavate, inflated, 15–25 µm long; conidia 12–23 × 10,5–15 µm,
with dark cap cells and 4 pendulous pale brown arms, 3–4-septate,
2,5–5 µm wide *C. parvulum*
2. Conidiophores ≤90 µm long; conidia 40–85 × 20–35 µm,
with 6–14 pendulous pale brown to yellowish brown or subhyaline arms,
15–20-septate, 2,5–4,5 µm wide *C. condensatum*
2. Conidiophores ≤45 µm long; conidia 20–45 × 12–30 µm,
with 3–9 pendulous subhyaline or pale brown arms,
3–9-septate, 4–6 µm wide *C. rilstonei*

Acknowledgments

The authors express their sincere gratitude to Prof. Bryce Kendrick and Dr. De-Wei Li for their critical review of the manuscript. The authors are grateful to Dr. P. M. Kirk for his taxonomic commentaries and opinion. The authors are grateful to the CNPq (proc. 141475/2013-7) for financial support and the “Programa de Pós-graduação em Botânica/UEFS” and ICMBIO for permission to collected microfungi in the “Floresta Nacional de São Francisco de Paula” (proc. 42334-1). The authors acknowledge the support provided by “Programa Ciência sem Fronteiras”. RFCR is grateful to the Cuban Ministry of Agriculture and “Programa de Salud Animal y Vegetal” (project P131LH003033) for facilities. We acknowledge the assistance provided by Dr. P.M. Kirk and Drs. V. Robert and A. Decock through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell’s editorial and Dr. Shaun Pennycook’s nomenclatural reviews are greatly appreciated.

Literature cited

- Castañeda-Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183, in: Anais do V Congresso Latino Americano de Micología. Brasília.
- Ellis MB. 1971. Dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey. [http://dx.doi.org/10.1016/S0181-1584\(01\)01057-0](http://dx.doi.org/10.1016/S0181-1584(01)01057-0)
- Ellis MB. 1976. More dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey. <http://dx.doi.org/10.5248/109.221>
- Schoknecht JD, Crane JL. 1977. Revision of *Torula* and *Hormiscium* species. *Torula occulta*, *T. diversa*, *T. elasticae*, *T. bigemina* and *Hormiscium condensatum* reexamined. *Mycologia* 69: 533–546. <http://dx.doi.org/10.2307/3758557>